

Protecting your baby from low blood glucose

Patient Information

Maternity Department



The Patient Information Leaflets page on the Trust website is available on the link:
<https://www.wwl.nhs.uk/patient-information-leaflets> or scan the QR code.

Author ID:	KA
Leaflet ref:	Obs 056
Version:	4
Leaflet title:	Protecting your baby from low blood glucose
Last review:	August 2022
Expiry date:	August 2024



What is low blood glucose?

You have been given this leaflet because your baby is at increased risk of having low blood glucose (also called low blood sugar or hypoglycaemia)

Babies who are small, premature, unwell at birth or whose mothers are diabetic or have taken certain medication (beta-blockers), may have low blood glucose in the first few hours and days after birth, and it is especially important for these babies to keep warm and feed as often as possible to maintain normal blood glucose levels.

If your baby is in one of these “at risk” groups, it is recommended that they have some blood tests to check their blood glucose level. Extremely low blood glucose, if not treated, can cause brain injury resulting in developmental problems. If low blood sugar is identified quickly, it can be treated to avoid harm to your baby.

What is blood glucose testing?

Your baby’s blood glucose is tested by a heel-prick blood test. A very small amount of blood is needed, and it can be done while you are holding your baby in skin-to-skin contact. The first blood glucose should be done before the start of the second feed (2-4 hours) after birth, and repeated until the blood glucose levels are stable.

You and your baby will need to stay in hospital for the blood tests.

You will know the result of the test straight away.

How can I avoid low blood glucose?

- **Skin-to-skin contact**

Skin to skin contact with your baby on your chest helps keep your baby calm and warm and helps establish breastfeeding. During skin-to-skin contact, your baby should wear a hat and be kept warm with a blanket or towel.

- **Keep your baby warm**

Put a hat on your baby for the first few days he/she is in hospital. Keep your baby in skin contact in your chest covered with a blanket and look into your baby’s eyes to check his/her well-being or keep warm with blankets if left in a cot.

- **Feed as soon as possible after birth**

Ask a member of staff to support you with feeding until you are confident and make sure you know how to tell if breastfeeding is going well, or how much formula to give your baby.

- **Feed as often as possible in the first few days**

Whenever you notice ‘feeding cues’ which include rapid eye movements under the eyelids, mouth and tongue movements, body movements and sounds or sucking on a fist, offer your baby a feed. Do not wait for your baby to cry – this can be a late sign of hunger.

- **Feed for as long as, or as much as, your baby wants.**

This is to ensure your baby gets as much milk as possible

- **Feed as often as baby wants, but do not leave your baby more than 3 hours between feeds**

If your baby is not showing any feeding cues yet, hold him/her skin-to-skin and start to offer a feed about 3 hours after the start of the previous feed.

- **Express your milk (colostrum)**

If you are breastfeeding and your baby struggles to feed, try to give some expressed breast milk. A member of staff will show you how to hand express your milk or watch the UNICEF hand expression video (search “UNICEF hand expression”) If possible it is good to have a small amount of expressed breast milk saved in case you need it later, so try to express a little breast milk in between feeds. Ask your midwife how to store your expressed milk.

What if I am worried about my baby?

Do not hesitate to speak to staff if you are worried. If your baby appears to be unwell, this could be a sign that they have a low blood glucose. As well as doing blood a test, staff will observe your baby to check he/she is well, but your observations are also important, as you are with your baby all the time so know your baby best. **It is important that you tell staff if you are worried**, that there is something wrong with your baby, as parents’ instincts are often correct.

How do I know if my baby is well?

- **Is your baby feeding well?**

In the first few days your baby should feed effectively at least 3 hours, until blood glucose is stable, and then at least 8 times in 24 hours. Ask a member of staff how to tell if your baby is attached and feeding effectively at the breast, or how much formula he/she needs. If your baby becomes less interested in feeding than before, this may be a sign they are unwell, and you should raise this with a member of staff.

- **Is your baby warm enough?**

Your baby should feel slightly warm to touch, although hands and feet can sometimes feel a little cooler. If you use a thermometer the temperature should be between 36.5°C and 37.5°C inclusive.

- **Is your baby alert and responding to you?**

When your baby is awake, he/she will look at you and pay attention to your voice and gestures. If you try to wake your baby, they should respond to you in the same way,

- **Is your baby’s muscle tone normal?**

A sleeping baby is very relaxed but should still have some muscle tone in their body, arms and legs and should respond to your touch. If your baby feels completely floppy, with no muscle tone when you lift their arms or legs, or if your baby is making strong repeated jerky movements, this is a sign they may be unwell. It can be normal to make brief, light jerky movements. Ask a member of staff if you are not sure about your baby’s movements.

- **Is your baby's colour normal?**

Look at the colour of the lips and tongue – they should be pink.

- **Is your baby breathing easily?**

A baby's breathing can be quite irregular, sometimes pausing for a few seconds and then breathing very fast for a few seconds. If you notice your baby is breathing very fast for a continuous period (more than 60 breaths per minute) or seems to be struggling to breathe with very deep chest movements, nostril flaring or making noises with each breath out – this is not normal.

What happens if my baby's blood glucose is low?

If the blood glucose test is low, your baby should feed as soon as possible and provide skin to skin contact. If the level is very low the neonatal team may advise urgent treatment to raise the blood glucose, and this could require immediate transfer to the Neonatal Unit. Another blood glucose test will be done before the next feed or within 2-4 hours.

If you are breastfeeding and your baby does not breastfeed straight away, a member of staff will review your baby to work out why. If he/she is happy that your baby is well he/she will support you to hand express your milk and give it by oral syringe or cup. If your baby has not breastfed, and you are unable to express any milk, you will be advised to offer formula.

In some hospitals the team may prescribe a dose of dextrose gel (sugar) as part of the feeding plan because this can be an effective way to bring your baby's glucose level up.

If you are breastfeeding and advised to give some infant formula, this is most likely to be for one or a few feeds only. You should continue to offer breastfeeds and try to express milk as often as possible to ensure your milk supply is stimulated.

Very occasionally, if babies are too sleepy or unwell to feed, or if the blood glucose is still too low after feeding, he/she may need to go to the Neonatal Unit. Staff will explain any treatment that might be needed. In most cases, low blood glucose quickly improves within 24-48 hours and your baby will have no further problems.

When can I go home with my baby?

It is recommended that your baby stays in hospital for 24 hours after birth. After that, if your baby's blood glucose is stable and he/she is feeding well, you will be able to go home.

Before you go home, make sure you know how to tell if your baby is getting enough milk. A member of staff will explain the normal pattern of changes in the colour of dirty nappies and number of wet nappies/dirty nappies. For further information, if you are breastfeeding, see 'How you and your midwife can recognise that your baby is feeding well' (Search 'UNICEF Baby Friendly assessment tool')

It is important to make sure that your baby feeds well **at least 8 times every 24 hours** and most babies feed more often than this.

There is no need to continue waking your baby to feed every 2-3 hours as long as he/she has had at least 8 feeds over 24 hours unless this has been recommended for a particular reason. You can now start to feed your baby responsively. Your midwife will explain this.

If you are bottle feeding, make sure you are not overfeeding your baby. Offer the bottle when he/she shows feeding cues and observe for signs that he/she wants a break. Don't necessarily expect your baby to finish a bottle – let him/her take as much milk as he/she wants.

Once you are home no special care is needed. As with all new-born babies, you should continue to look for signs that your baby is well and seek medical advice if you are worried at all about your baby.

Further Information and References

British Association of Perinatal Medicine Identification and Management of Neonatal Hypoglycaemia in the Full Term Infant – A Framework for Practice 2017.

UNICEF: www.unicef.org.uk/BabyFriendly

Comments, Compliments or Complaints

The Patient Relations/Patient Advice and Liaison Service (PALS) Department provides confidential on the spot advice, information and support to patients, relatives, friends, and carers.

Contact Us

Tel: 01942 822376 (Monday to Friday 9am until 4pm)

The Patient Relations/PALS Manager
Wrightington, Wigan and Leigh Teaching Hospitals NHS Foundation Trust
Royal Albert Edward Infirmary
Wigan Lane
Wigan
WN1 2NN

Ask 3 Questions

Become more involved in decisions about your healthcare. You may be asked to make choices about your treatment. To begin with, try to make sure you get the answers to three key questions:

1. What are my options?
2. What are the positives and negatives of each option for me?
3. How do I get support to help me make a decision that is right for me?



How We Use Your Information

For details on how we collect, use, and store the information we hold about you, please see patient information leaflet, Ref. **Corp 006** How we use your information, this can be found on the Patient Information Leaflets page on the Trust website, see details on the front cover.

This leaflet is also available in audio, large print, Braille, and other languages upon request. For more information, please ask in the department/ward.

© Wrightington, Wigan and Leigh Teaching Hospitals NHS Foundation Trust.
All rights reserved. Not to be reproduced in whole or in part without the permission of the copyright owner.

Call 111 first when it's less urgent than 999.



EMPLOYER RECOGNITION SCHEME

GOLD AWARD 2021

Proudly serving those who serve.

Phone: 0808 802 1212

Text: 81212

www.veteransgateway.org.uk

