

Female Sterilisation

Patient Information

Obstetrics & Gynaecology Department



The Patient Information Leaflets page on the Trust website is available on the link:
<https://www.wvl.nhs.uk/patient-information-leaflets> or scan the QR code.

Author ID: JAD
Leaflet ref: Gyn 006
Version: 8
Leaflet title: Female Sterilisation
Last review: October 2024
Expiry Date: October 2026

Introduction

Sterilisation is a permanent method of contraception that should only be considered by those who are absolutely sure that they will never want any further children. In women, the fallopian tubes are blocked, cut, or removed by means of an operation, so that the sperm and the eggs will not meet and fertilise thereby causing a pregnancy.

Female sterilisation is not always as reliable as other methods used correctly, especially male sterilisation, which is a safer and more effective operation.

Alternative methods of Contraception

Method	Number of pregnancies per 1,000 women using method correctly for 1 year
Combined Contraceptive Pill	15 (2 – 30)
Progesterone only pill (mini pill)	35 (14 – 43)
Copper IUCD (coil)	3 (1 – 5)
Progesterone IUCD (Mirena coil)	1 (0 – 2)
Depo-Provera (pill injection)	3 (0 – 10)
Implanon (implant)	0.5 (0 – 0.5)
Male Sterilisation (Vasectomy)	1 (0 – 2)
Female Sterilisation	2 (0 – 5)

Sterilisation is the only method which is permanent.

Female sterilisation does not make periods either better or worse and if you have period problems other methods of contraception may be more suitable.

Method of performing a Sterilisation

Laparoscopic (Operation in theatre with a general anaesthetic)

In order to perform this type of sterilisation it is necessary to admit you for the day to Leigh Infirmary. You will be given a general anaesthetic. The sterilisation is performed by laparoscopy (see separate sheet for details) during which clips can be applied to the tubes to block them or the whole tube can be removed. Laparoscopic (keyhole) sterilisation is, in the majority of cases, a safe simple procedure with no complications but, as with any operation, complications can arise. This occurs in about 2 cases out of every 1000 laparoscopic sterilisations. Possible complications are bleeding or injury to the bladder or bowel when the instruments are inserted.

In some cases, the complication can be sorted out without any further incisions but occasionally it is necessary to perform an open operation. It is important to realise that, although complications are rare, they can occasionally be fatal; for every 12,000 sterilisations performed one lady will die (0.00008%).

Reversal of a Sterilisation

A sterilisation operation should be regarded as permanent. Although reversal of a clip sterilisation can be attempted, success rates are often low, and the procedure is not offered under the NHS. If the tubes are completely removed reversal is never possible.

Failure of Sterilisation

Even if the procedure is performed entirely correctly pregnancies can occur after sterilisation and for clip sterilisation this happens in about 1 in 200 cases for the laparoscopic method compared with 1 in 2000 for vasectomy. It can occur at any time after sterilisation, even many years later.

Failure of sterilisation is more common, even if the tubes are completely removed, if you are or have recently been pregnant at the time of the sterilisation. For this reason, combined with the fact that you are more likely to regret the procedure if it is done during delivery or close to the time of the pregnancy, it is usually better to return for it to be done about 3 months later.

If a pregnancy does occur following sterilisation there is an increased chance that it will be an ectopic pregnancy (pregnancy in the tube) since the tube will have been damaged by the sterilisation procedure and may therefore trap the egg inside the tube. As this can be a dangerous condition it is very important that you seek advice immediately if you suspect that you are pregnant after sterilisation.

Even if the whole tube is removed an ectopic pregnancy can still occur in the cut end of the tube.

Female sterilisation is not always as reliable as other methods used correctly, especially male sterilisation, which is a safer and more effective operation.

Contraception around the time of Sterilisation

A sterilisation operation will not prevent a pregnancy that has already occurred. Sterilisation is contraceptive because it stops the eggs passing along the tubes to the uterus. If an egg has just been released before the tubes are blocked or removed it could still become fertilized. It is therefore very important that you do continue using contraception until the next period after a laparoscopic sterilisation operation.

Comments, Compliments or Complaints

The Patient Relations/Patient Advice and Liaison Service (PALS) Department provides confidential on the spot advice, information and support to patients, relatives, friends, and carers.

Contact Us

Tel: 01942 822376 (Monday to Friday 9am to 4pm)

The Patient Relations/PALS Manager
Wrightington, Wigan and Leigh Teaching Hospitals NHS Foundation Trust
Royal Albert Edward Infirmary
Wigan Lane
Wigan WN1 2NN

Ask 3 Questions

Become more involved in decisions about your healthcare. You may be asked to make choices about your treatment. To begin with, try to make sure you get the answers to three key questions:

1. What are my options?
2. What are the pros and cons of each option for me?
3. How do I get support to help me make a decision that is right for me?



How We Use Your Information

For details on how we collect, use, and store the information we hold about you, please see patient information leaflet, Ref. **Corp 006** How we use your information, this can be found on the Patient Information Leaflets page on the Trust website, see details on the front cover.

This leaflet is also available in audio, large print, Braille, and other languages upon request.

For more information, please ask in the department/ward.

© Wrightington, Wigan and Leigh Teaching Hospitals NHS Foundation Trust.
All rights reserved. Not to be reproduced in whole or in part without the permission of the copyright owner.

Call 111 first when it's less urgent than 999.



EMPLOYER RECOGNITION SCHEME

SILVER AWARD

Proudly supporting those who serve.

Phone: 0808 802 1212

Text: 81212

www.veteransgateway.org.uk

