

Exercises for base of thumb pain

Patient Information

Hand Therapy Team



The Patient Information Leaflets page on the Trust website is available on the link:
<https://www.wwl.nhs.uk/patient-information-leaflets> or scan the QR code.

Author ID: AP
Leaflet Ref: OT 015
Version: 1
Leaflet title: Exercises for base of thumb pain.
Date Produced: May 2024
Expiry Date: May 2026

Exercise

Exercise can help to reduce joint stiffness, improve flexibility, and reduce pain. Exercise can strengthen the muscles around the base of the thumb and within the hand to support the function of the thumb. Your therapist will advise you on the frequency and repetition of each exercise. Sometimes, exercising the thumb can be uncomfortable. If the exercises generate pain, this should last no longer than 24 hours before recovery. Exercise should not be repeated until your pain has resolved and may require a rest period before re-commencing. If discomfort persists for more than 24 hours after exercising, contact your therapist for advice.


Warming up

Heat and warmth can be beneficial to movement and stretch and is an important preparation for exercise. Warm the hand in a bowl of warm water for 2-3 minutes. Move your thumb in a circular, clockwise direction within the water for about a minute. After this time, rotate the thumb in an anti-clockwise direction within the water for a further minute.

Stretches


Sometimes the muscles around the thumb can become tight and the thumb cannot stretch away from the palm as wide as it used to. This is called the thumb web space, and it can tighten in patients with thumb Osteoarthritis (OA). This can limit movement of the thumb and cause poor positioning of the thumb in activity.

If it is appropriate, your therapist will show you techniques to relieve pain, improve movement and stretch any tightness of the soft tissues. Your therapist will advise you individually on what is best for your thumb. These techniques may be introduced before certain exercises to put your thumb in a better position before movement.



Positioning

When performing exercises of the thumb, it is important to be aware of your thumb's positioning. The joints of your thumb work best in function when they are slightly bent. Maintaining a wide web space and stable thumb position when pinching, known as an 'O' shape pinch, is the best way to use your thumb in functional tasks.



Exercise Programme


These exercises are given in a sequence, and you should progress only when the first stage exercises can be performed well. Your therapist can guide you on your progression. It is important to complete the exercises accurately to ensure correct thumb positioning. Movements of the thumb should be performed slowly and under good control.

Stage 1

1a) Thumb Extension


Rest your hand on a table on the little finger side, with the thumb relaxed.

Lift your thumb upwards, keeping the end thumb joint slightly bent (as shown)




Hold this position for 10 seconds and then relax. Repeat the exercise 10 times.

Do not over straighten (hyperextend) the joints of the thumb (as shown).



1b) Thumb Abduction

Rest your hand on a table on the little finger side, with the thumb relaxed. Lift your thumb outwards, keeping the end thumb joint slightly bent (as shown).




Hold this position for 10 seconds and then relax. Repeat the exercise 10 times.

1c) Opposition

Rest your hand on a table on the little finger side, with the thumb relaxed.

Bring your thumb across to touch the tip of each finger. Ensure you make an 'O' shape when the fingertips touch, keeping the thumb joints slightly bent (as shown).

Hold this position for 10 seconds and then relax. Repeat the exercise 10 times.




Do not pinch through the thumb with the end joint in a straight or hyperextended position (as shown).





1d) Proprioception

Place a tennis ball in your palm with your fingers placed over the top and your thumb relaxed to the side (as shown)

Stretch your thumb outwards around the ball and then rotate the ball around using your fingers and thumb.



Following the white line on the tennis ball can be useful to focus your finger and thumb movements.





Try to keep the thumb in an 'O' shape posture whilst moving the ball around.
Continue for 2 -3 minutes.

Stage 2

2a) Resisted Extension

Place a rubber band or equivalent around your hand and thumb (as shown).

The band should go across the middle section of the thumb.




Stretch the band slowly upwards. Try to keep the joints of the thumb slightly bent when pulling against the band.

Hold the stretch for 5 seconds against the pull of the band, and then slowly release. It is equally important to control the recoil of the band when relaxing the thumb; do not let the band pull the thumb down.

Repeat 10 times.


Be careful not to over straighten the thumb joints (as shown).



2b) Resisted Abduction

Place a rubber band or equivalent around your hand and thumb (as shown).

The band should go across the middle section of the thumb.



Stretch the band outwards whilst keeping the end joints of the thumb slightly bent.


Hold the stretch for 5 seconds against the pull of the band, and then slowly release. It is equally important to control the recoil of the band when relaxing the thumb, do not let the band pull the thumb back in.

Repeat 10 times.

2c) Resisted finger abduction

Place a rubber band or equivalent around your fingers (as shown).

Stretch the index finger outwards against the resistance of the band. Hold the stretch for 5 seconds against the pull of the band, and then slowly release. It is equally important to control the recoil of the band when relaxing the index finger; do not let the band pull the finger back.




Repeat 5 times.

Progress the exercise by bringing your index finger and thumb together into an 'O' shape posture, pulling against the resistance of the band.

Hold the stretch for 5 seconds against the pull of the band, and then slowly release. It is equally important to control the recoil of the band when relaxing the index finger; do not let the band pull the finger back.

Repeat 10 times.



Stage 3



Pinch postures

In this final stage, the position of your thumb in function is looked at. The posture your thumb adopts when pinching is important for stable loading through the base of the thumb.

Keeping the joints of the thumb slightly bent when applying pinch, and maintaining an 'O' shape pinch posture, is the best position to use the thumb in.

3a) Pen grip



Practise writing a sentence using the correct thumb posture. Repeat the sentence 5 times.



3b) Paper tear

Practice tearing a piece of paper using the correct thumb position (as shown).



Repeat this 10 times, maintaining a good posture of the thumb as you tear.



3c) Peg pinch

Practice gently squeezing a peg using the 'O' shape pinch (as shown).

Squeeze the peg and then slowly release the pinch, whilst maintaining the correct thumb posture





3d) Key turn

Practice holding a key using the correct thumb position (as shown).

Place the key in the lock and turn, whilst maintaining the correct thumb posture.

Repeat this 10 times.



Once you can maintain the correct thumb posture, try to use it consistently in all functional activities when pinching or gripping with the thumb. New patterns of movement take time to become established in the brain, so repetition is important to encourage this to become the new norm.

Disclaimer

Not all exercises are suitable for everyone, and while every precaution has been taken in selecting these exercises, they do not come without risks. If you have any concerns regarding these exercises, discuss them with your therapist.

The exercise plans shown in this leaflet are designed to be used as a guide and a point of reference; they can be modified and progressed under the supervision of your therapist.

Comments, Compliments or Complaints

The Patient Relations/Patient Advice and Liaison Service (PALS) Department provides confidential on the spot advice, information and support to patients, relatives, friends, and carers.

Contact Us

Tel: 01942 822376 (Monday to Friday 9am until 4pm)

The Patient Relations/PALS Manager
Wrightington, Wigan and Leigh Teaching Hospitals NHS Foundation Trust
Royal Albert Edward Infirmary
Wigan Lane
Wigan
WN1 2NN

Ask 3 Questions

Become more involved in decisions about your healthcare. You may be asked to make choices about your treatment. To begin with, try to make sure you get the answers to three key questions:

1. What are my options?
2. What are the positives and negatives of each option for me?
3. How do I get support to help me make a decision that is right for me?



How We Use Your Information

For details on how we collect, use, and store the information we hold about you, please see patient information leaflet, Ref. **Corp 006** How we use your information, this can be found on the Patient Information Leaflets page on the Trust website, see details on the front cover.

This leaflet is also available in audio, large print, Braille, and other languages upon request. For more information, please ask in the department/ward.

© Wrightington, Wigan and Leigh Teaching Hospitals NHS Foundation Trust.
All rights reserved. Not to be reproduced in whole or in part without the permission of the copyright owner.

Call 111 first when it's less urgent than 999.



EMPLOYER RECOGNITION SCHEME

GOLD AWARD 2021

Proudly serving those who serve.

Phone: 0808 802 1212

Text: 81212

www.veteransgateway.org.uk

